

Please enter first then last names	YYYY-MM-DD
Patient Name:	Patient Birthdate:
HCP Evaluator:	Evaluation Date:
<p>Scoring System: 0 = unable to evaluate or absent: 1 = mild or intermittent: 2 = severe Ratings should be based on symptoms and signs occurring during the week prior to interview. No score should be given if symptoms result from physical disability or illness.</p>	
Please check the score that applies with an X	<p>NOTE: The scale is designed as a screening tool and is not diagnostic</p>
0 1 2	<p>A. Mood-Related Signs Anxiety: anxious expression, ruminations, worrying Sadness sad expression, sad voice, tearfulness Lack of reactivity to pleasant events Irritability: easily annoyed, short-tempered</p>
0 1 2	<p>A. Behavioral Disturbance Agitation: restlessness, handwringing, hairpulling Retardation: slow movements, slow speech, slow reactions Multiple physical complaints (score 0 if GI symptoms only) Loss of interest a 0 1 2 less involved in usual activities (score only if change occurs acutely: i.e., in less than one month)</p>
0 1 2	<p>A. Physical Signs Appetite loss: eating less than usual Weight loss: score 2 if greater than 5 lb. in one month Lack of energy: fatigues easily, unable to sustain activities (score only if change occurred acutely, i.e., in less than 1 month)</p>
0 1 2	<p>A. Cyclic Functions Diurnal variation of mood: symptoms worse in the morning Difficulty falling asleep: later than usual for this individual Multiple awakenings during sleep Early-morning awakening: earlier than usual for this individual</p>
0 1 2	<p>A. Ideational Disturbance Suicide: feels life is not worth living, has suicidal wishes or makes suicide attempt Poor self-esteem a 0 1 2 self-blame, self-deprecation, feelings of failure Pessimism: anticipation of the worst Mood-congruent delusions: delusions of poverty, illness or loss</p>
Score:	<p>A score >10 probably major depressive episode. A score >18 definite major depressive episode</p>